

Recocam

20mg/ml Meloxicam Solution for Injection

INDICATIONS

Horses:

- For use in the alleviation of inflammation and pain in both acute and chronic musculo-skeletal disorders
- For the relief of pain associated with equine colic

Cattle:

- For use in acute respiratory infections with appropriate antibiotic to reduce signs in cattle
- For use in diarrhoea in combination with oral re-hydration therapy to reduce clinical signs in calves over one week of age and young non-lactating cattle
- For adjunctive therapy in the treatment of acute mastitis in combination with antibiotic therapy

Pigs:

- For use in non-infectious locomotor disorders to reduce the symptoms of lameness and inflammation
- For adjunctive therapy in the treatment of puerperal septicaemia and toxæmia (mastitis-metritis-agalactia syndrome) with appropriate antibiotic therapy



BENEFITS

- Convenient: Single injection course in cattle
- COX -2 inhibitory mode of action
- Multi-species use
- Multi-indication use
- Licensed for use in diarrhetic calves
- IV & SC cattle option

LIST NO.	UNIT	CASE
1REC001	100ml	12



See reverse side for Administration and Dosage.

Practice support line: 0800 6524463

Recocam

20mg/ml Meloxicam Solution for Injection

PRESENTATION

A clear yellowish solution containing 20mg/ml meloxicam for injection.

TARGET SPECIES

Horses, Cattle and Pigs

USES

Horses:

For use in the alleviation of inflammation and relief of pain in both acute and chronic musculo-skeletal disorders. For the relief of pain associated with equine colic.

Cattle:

For use in acute respiratory infection with appropriate antimicrobial therapy to reduce clinical signs in cattle. For use in diarrhoea in combination with oral rehydration therapy to reduce clinical signs in calves of over one week of age and young, non-lactating cattle. For adjunctive therapy in the treatment of acute mastitis, in combination with antimicrobial therapy.

Pigs:

For use in non-infectious locomotor disorders to reduce the symptoms of lameness and inflammation. For adjunctive therapy in the treatment of puerperal septicaemia and toxæmia (mastitis –metritis –agalactia syndrome) with appropriate antimicrobial therapy.

AMOUNTS TO BE ADMINISTERED AND ADMINISTRATION ROUTE

Horses:

Single intravenous injection at a dosage of 0.6mg meloxicam/kg bodyweight (i.e. 3.0ml/100kg bodyweight).

For use in the alleviation of inflammation and relief of pain in both acute and chronic musculo-skeletal disorders. Oral suspensions of Recocam may be used for continuation of treatment at a dosage of 0.6mg meloxicam/kg bodyweight 24 hours after the administration of the injection.

Avoid introduction of contamination during use. Do not broach the stopper more than fifty times.

Cattle:

Single subcutaneous or intravenous injection at a dosage of 0.5mg meloxicam/kg bodyweight (i.e. 2.5ml/100kg bodyweight) in combination

with antimicrobial therapy or with oral rehydration therapy, as appropriate.

Pigs:

Single intramuscular injection at a dosage of 0.4mg meloxicam/kg bodyweight (i.e. 2.0ml/100kg bodyweight) in combination with antimicrobial therapy, as appropriate. If required, a second administration of Recocam can be given after 24 hours.

CONTRA-INDICATIONS & WARNINGS

i) Special precautions for use in animals

- Do not use in horses less than 6 weeks of age.
- Do not use in animals suffering from impaired hepatic, cardiac or renal function and haemorrhagic disorders, or where there is evidence of ulcerogenic gastrointestinal lesions.
- Do not use in case of hypersensitivity to the active substance or to any of the excipients.
- For the treatment of diarrhoea in cattle, do not use in animals of less than one week of age.
- If adverse reactions occur, treatment should be discontinued and the advice of a veterinarian should be sought.
- Avoid use in very severely dehydrated, hypovolaemic or hypotensive animals which require parenteral rehydration, as there may be a potential risk of renal toxicity.
- In case of inadequate relief of pain when used in the treatment of equine colic, careful re-evaluation of the diagnosis should be made as this could indicate the need for surgical intervention.

ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals

Accidental self-injection may give rise to pain. People with known hypersensitivity to non-steroidal anti-inflammatory drugs (NSAIDs) should avoid contact with the veterinary medicinal product. In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

ADVERSE REACTIONS (FREQUENCY AND SERIOUSNESS)

In horses, a transient swelling at the injection site can occur but resolves without intervention. In very rare cases anaphylactoid reactions may occur and should be treated symptomatically.

In cattle, subcutaneous and intravenous administration is well tolerated; only a slight transient swelling at the injection site following subcutaneous administration was observed in less than 10% of the cattle treated in clinical studies.

In pigs, subcutaneous intramuscular administration is well tolerated.

Interaction with other medicinal products and other forms of interaction

Do not administer concurrently with glucocorticosteroids, other non-steroidal anti-inflammatory drugs or with anti-coagulant agents.

USE DURING PREGNANCY AND LACTATION

Horses:

Do not use in pregnant or lactating mares. Do not use in horses producing milk for human consumption.

Cattle and Pigs:

Can be used during pregnancy and lactation.

WITHDRAWAL PERIOD(S)

Horses.

Meat and Offal: 5 days.

Not authorised for horses producing milk for human consumption.

Cattle.

Meat and Offal: 15 days.

Milk: 5 days.

Pigs.

Meat and Offal: 5 days.

LEGAL CATEGORY

POM-V

MARKETING AUTHORISATION HOLDER

Bimeda Chemicals Ltd., Broomhill Road, Tallaght, Dublin 24. Ireland.

MARKETING AUTHORISATION NUMBER

EU/2/11/133/001-003

Advice should be sought From Medicine Prescriber before using Recocam.

Contact: Bimeda Inc, Unit 2, Bryn Cefni Industrial Park, Llangefni, Anglesey, Wales.

TAKE TIME



OBSERVE LABEL DIRECTIONS

Use Medicines Responsibly
Advice should be sought from a prescriber before using the product
www.noah.co.uk

